

Package leaflet: Information for the patient

STANOZOLOL ® 10 microgram Tablets

Read all of this leaflet carefully before you start taking this medicine because it contains Important information for you.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it onto others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible Side effects not listed in this leaflet.

What is in this leaflet

1. *What Stanozolol is and what it is used for*
2. *What you need to know before you take Stanozolol*
3. *How to take Stanozolol*
4. *Possible side effects*
5. *How to store Stanozolol*
6. *Other information*

1. WHAT STANOZOLOL IS AND WHAT IT IS USED FOR

Composition:

One tablet contains:

Active substance: 10 mg
Stanozolol, Magnesium stearate,
FD & C, Ludipress (lactose,
povidone, crospovidone)

Drug formulation:

Round 10 mg white and flat tablets.

Pharmacological properties:

Stanozolol is an anabolic steroid. It stimulates anabolic processes and inhibits catabolic effects caused by

glucocorticoids. This leads to an increase in muscle mass, reducing body fat levels, improves connective tissue. Androgenic (moderate) effect may contribute to the development of male secondary sexual signs.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE STANOZOLOL

Indications:

In disorders of protein synthesis, traumas, burns, post-operative infections, renal failure, osteoporosis, corticosteroid therapy, aplastic anemia
Treatment of tuberculosis,
Treatment of hepatitis, Treatment of asthma.

3. HOW TO TAKE STANOZOLOL

Dosage and route of administration:

Doses: Initial dose 10-20 mg per day. Maintenance dose: 5-10 mg

per day.

When we want to reduce the incidence of angioedema attacks, we need to take the dose individually. The starting dose is 2 mg three times a day

4. POSSIBLE SIDE EFFECTS

Side effects:

Women: virilizing symptoms, change in menstrual cycle, inhibition of ovarian function

Hypercalcemia

Men: before puberty - virilizing symptoms, pigmentation of the skin, post-puberty bladder inflammation, Priapizym, adults - prostate cancer.

Patients should be instructed to report any of the following: nausea, vomiting, changes in skin color, ankle swelling, too frequent or persistent erections of the penis.

Contraindications:

Drug hypersensitivity, prostate cancer, breast cancer, atherosclerosis, renal failure, pregnancy, breastfeeding

Interaction with other medicines:

Increases the effect of anticoagulants, antiplatelet agents, increases side effects of medicines.

5. HOW TO STORE STANOZOLOL

It should be stored in a dry place away from heat and humidity. 15° to 25° C (59° to 77° F)

Keep Stanozolol out of the reach of children and away from pets. Do not refrigerate or freeze. Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your Pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. OTHER INFORMATION

Expiration date:

3 years. Do not use after expiry date.

Terms of Use:

By doctor's prescription

MARKETING AUTHORISATION HOLDER

MANUFACTURER:

Canadian Pharmaceuticals
32 Tower Ave
Steinbach, MB R5G 0B8
Canada

This leaflet was updated in July 2009