

Package leaflet: Information for the patient
DECA DURABOLIN® 250 mg/ml, solution for injection

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it onto others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet

1. *What Deca Durabolin is and what it is used for*
2. *What you need to know before you take Deca Durabolin*
3. *How to take Deca Durabolin*
4. *Possible side effects*
5. *How to store Deca Durabolin*
6. *Other information*

1. WHAT DECA DURABOLIN IS AND WHAT IT IS USED FOR

Composition:

Each ml of the solution contains:
Nandrolone Decanoate - 250 mg,
Benzyl alcohol, Benzyl benzoate,
Arachis oil.

Form of the drug:

Clear oil solution from slightly yellow to yellow color with a characteristic smell

Pharmacological properties:

Synthetic production of testosterone with weak androgenicity and high anabolic efficacy

The anabolic effects activates processes in the epithelium bone and muscle tissue as a result of stimulating the synthesis of proteins and structural components of the cells. This enlarges the assimilation of amino acids from the

small intestines. And the leads to positive nitrogen balance. It increases LDL concentration and decrease HDL. These effects disappear once the treatment is stopped.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE DECA DURABOLIN

Indications:

Treating anemia, Treatment of renal impairment, breast cancer, Osteoporosis, Diabetic retinopathy, Full burns

3. HOW TO TAKE DECA DURABOLIN

Dosage and How to Take It:

This medicine is given intramuscularly as directed by your pharmacist. It should not be administered intravenously. Intramuscular injections should be given deep in the gluteal muscle.

Anemia - 100-200 mg (women - 50-100 mg) once a week;

For stimulating the anabolic processes - 50-200 mg (women - 50-100 mg) once a week (for one or four weeks).

Children from 2-13 years - 0.4 mg / kg. Once every three or four weeks. The treatment should be continue from 1-4

weeks between the injections. The treatment should be repeated over 8 weeks

With this treatment we need to keep a diet rich in protein.

IT SHOULD NOT BE GIVEN INTRAVENOUSLY!

4. POSSIBLE SIDE EFFECTS

Side effects:

Women: virilizing symptoms, change in the menstrual period, inhibition of ovarian function
Hypercalcemia, Menstrual cycle disorders

Men: before puberty – virilizing symptoms, Pigmentation of the skin, after puberty inflammation of the bladder, Priapizym. Adults - prostate cancer, Hypertrophy

Men and women: atherosclerosis, bleeding, Jaundice disease, Bone pain
Patients should be instructed to report any of the following: nausea, vomiting, changes in skin color, ankle swelling, too frequent or persistent erections of the penis.

Contraindications:

Oversensitivity to the drug, prostate cancer, breast cancer (women with hypercalcemia), atherosclerosis, kidney failure, pregnancy, Lactation

Overdose:

There have been no reports of acute overdose with androgens.

Interactions with other medicines:

Glucocorticoids, Mineralocorticoids, Corticotropin preparations, with sodium that increases fluid retention in the body, Increase the risk of swelling,

Acne.

Increases antiplatelet activity, Anticoagulants, Insulin, decrease the effect of Somatotropin and its derivatives.

5. HOW TO STORE DECA DURABOLIN

It should be stored in a dry place away from heat and humidity. 15° to 25° C (59° to 77° F)

Keep Deca Durabolin out of the reach of children and away from pets. Do not refrigerate or freeze;

Store in the original package in order to protect from light

Do not throw away any medicines via wastewater or household waste. Ask your Pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. OTHER INFORMATION

Expiry date:

5 years. Do not use after the expiry date stated on the pack.

Terms of Use:

By doctor's prescription

MARKETING AUTHORISATION HOLDER MANUFACTURER:

Canadian Pharmaceuticals
32 Tower Ave
Steinbach, MB R5G 0B8
Canada

This leaflet was updated in July 2006