

Package leaflet: Information for the patient  
**WINSTROL® 75 mg/ml solution for injection**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it onto others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

*What is in this leaflet*

1. *What Winstrol is and what it is used for*
2. *What you need to know before you take Winstrol*
3. *How to take Winstrol*
4. *Possible side effects*
5. *How to store Winstrol*
6. *Other information*

## **1. WHAT WINSTROL IS AND WHAT IT IS USED FOR**

Each milliliter of the oily solution contains the following: Active substance: 75 mg Stanozolol. The other ingredients are: The solution for injection contains Arachis oil and Benzyl Alcohol.

### **Form of the drug:**

Clear oil solution from slightly yellow to yellow colour with characteristic smell

### **Pharmacological properties:**

Winstrol is an anabolic steroid. It stimulates the anabolic processes and inhibits the catabolic ones, which is caused by glucocorticoids. That brings to an increase of the muscle mass, it postpones the fat distribution, it

improves the connective tissue. The androgenic effect (a moderate one) can contribute to the development of the male secondary sexual characteristics.

## **2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE WINSTROL**

### **Indications:**

Protein synthesis disorder, Traumas , Burns, Postoperative Infections, kidney failure, Osteoporosis, Corticosteroid therapy, Aplastic anemia, Negative nitrogen balance  
Treatment of tuberculosis, Treatment of hepatitis, Treatment of asthma, Treatment of Osteomyelitis

## **3. HOW TO TAKE WINSTROL**

### **Dosage and How to Take It:**

This medicine is given intramuscularly as directed by your pharmacist. It should not be administered intravenously. Intramuscular injections should be given deep in the gluteal muscle.

The dose depends from the patients, the disease state, sex, Clinical effectiveness.

Middle dose for men: 50-100 mg for a

day; women 25-50 mg for a week. .  
When we want to decrease the frequency of the angioedema attacks we should take the dose individually.  
IT SHOULD NOT BE GIVEN INTRAVENOUSLY!

#### **4. POSSIBLE SIDE EFFECTS**

##### **Side effects:**

Women: virilizing symptoms, change in the menstrual period, inhibition of ovarian function  
Hypercalcemia

Men: before puberty - virilizing symptoms , Pigmentation of the skin, after puberty inflammation of the bladder , Priapizym, adults - prostate cancer, Hypertrophy

Men and women: atherosclerosis , bleeding, Jaundice disease, Bone pain  
Patients should be instructed to report any of the following: nausea, vomiting, changes in skin color, ankle swelling, too frequent or persistent erections of the penis.

##### **Contraindications:**

Oversensitivity to the drug, prostate cancer, breast cancer, atherosclerosis, kidney failure, pregnancy, Breastfeeding

##### **Overdose:**

In the event of an overdose, medical advice should be sought

Interactions with other medicines:  
Increases the action of Anticoagulants, Antiplatelet agents, increases the side effects of the drugs.

#### **5 HOW TO STORE WINSTROL.**

It should be stored in a dry place away from heat and humidity. 15° to 25° C (59° to 77° F)

Keep Winstrol out of the reach of children and away from pets. Do not refrigerate or freeze

Store in the original package in order to protect from light

Do not throw away any medicines via wastewater or household waste. Ask your Pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

#### **6. OTHER INFORMATION**

##### **Expiration date:**

3 years. Do not use after expiry date.

##### **Terms of Use:**

By doctor's prescription

**MARKETING AUTHORITY**  
**HOLDER**

##### **MANUFACTURER:**

Canadian Pharmaceuticals  
32 Tower Ave  
Steinbach, MB R5G 0B8  
Canada

*This leaflet was updated in July 2009*

